

## Author Index

Abuin, G., Colli, W., De Souza, W. and Alves, M.J.M.  
A surface antigen of *Trypanosoma cruzi* involved in cell invasion (Tc-85) is heterogeneous in expression and molecular constitution, 229

Akella, R., See Vaidya, A.B., 97

Allsopp, B., Carrington, M., Baylis, H., Sohal, S., Dolan, T. and Iams, K.  
Improved characterization of *Theileria parva* isolates using the polymerase chain reaction and oligonucleotide probes, 137

Alunda, J.M., see Balaña-Fouce, R., 43

Alves, M.J.M., see Abuin, G., 229

Asikin, N., see Grandea III, A.G., 31

Atiya, A., see Zidovetzki, R., 199

Balaña-Fouce, R., Ordóñez, D. and Alunda, J.M.  
Putrescine transport system in *Leishmania infantum* promastigotes, 43

Banerjee, A.B., see Ghoshal, K., 21

Baylis, H., see Allsopp, B., 137

Beach, D.H., Pascal, Jr., R.A. and Holz Jr., GG.  
Effects of thiastearyl acids on growth and on dihydrostearic acid and other phospholipid fatty acyl groups of *Leishmania* promastigotes, 57

Beecroft, R.P., see Tolson, D.L., 109

Boyd, D., De Vos, T., Klassen, G. and Dick, T.  
Characterization of the ribosomal DNA from *Trichinella spiralis*, 67

Breier-Saraiva, E.M., see Silva Filho, F.C., 73

Carrington, M., see Allsopp, B., 137

Casper, P., Gentz, R., Matile, H., Pink, J.R. and Sinigaglia, F.  
The circumsporozoite protein gene from NF54, a *Plasmodium falciparum* isolate used in malaria vaccine trials (*Short Communication*), 185

Chambers, A.E., see Harnett, W., 119

Chaudhuri, J. and Donahue, M.J.  
Serotonin receptors in the tissues of adult *Ascaris suum*, 191

Cohen, C., see Grandea III, A.G., 31

Colli, W., see Abuin, G., 229

Dalton, J.P. and Heffernan, M.  
Thiol proteases released in vitro by *Fasciola hepatica*, 161

Davis, T.B., see Grandea, III, A.G., 31

Davis, W.C., see Reduker, D.W., 239

De Boeck, H., see Zidovetzki, R., 199

De Souza, W., see Abuin, G., 229

De Souza, W., see Silva Filho, F.C., 73

De Souza, W., see Lafaille, J.J., 127

De Vos, T., see Boyd, D., 67

Dick, T., see Boyd, D., 67

Dolan, T., see Allsopp, B., 137

Donahue, M.J., see Chaudhuri, J., 191

Dresden, M.H., see Rege, A.A., 89

Fidge, N.H., see Rogers, M.V., 79

Fuhrman, J.A. and Piessens, W.F.  
A stage-specific calcium-binding protein from microfilariae of *Brugia malayi* (Filariidae), 249

Gentz, R., see Caspers, P., 185

Gero, A.M.  
Induction of nucleoside transport sites into the host cell membrane of *Babesia bovis* infected erythrocytes, 269

Ghoshal, K., Banerjee, A.B. and Ray, S.  
Methylglyoxal-catabolizing enzymes of *Leishmania donovani* promastigotes, 21

Goff, W.L., see Reduker, D.W., 239

Goldenberg, S., see Lafaille, J.J., 127

Grande III, A.G., Tuyen, L.K., Asikin, N., Davis, T.B., Philipp, M., Cohen, C. and McReynolds, L.A.  
A  $\lambda$ gt11 cDNA recombinant that encodes *Dirofilaria immitis* paramyosin, 31

Hannon, G.J., see Maroney, P.A., 277

Harnett, W., Chambers, A.E., Renz, A. and Parkhouse, R.M.E.  
An oligonucleotide probe specific for *Onchocerca volvulus*, 119

Hedstrom, L. and Wang, C.C.  
Purine base transport in wild-type and mycophenolic acid-resistant *Trichomonas foetus*, 219

Heffernan, M., see Dalton, J.P., 161

Henkle, K.J., see Rogers, M.V., 79

Herrera, P.R., see Rege, A.A., 89

Higashi, G.I., see Werner, C., 209

Holz, Jr., G.G., see Beach, D.H., 57

Howard, R.J., see Panton, L.J., 149

Iams, K., see Allsopp, B., 137

Jansen, J., see Wesseling, J.G., 167

Jasmer, D.P., see Reduker, D.W., 239

Klassen, G., see Boyd, D., 67

Krieger, M.A., see Lafaille, J.J., 127

Kwakye-Berko, F. and Meshnick, S.R.  
Binding of chloroquine to DNA, 51

Lafaille, J.J., Linss, J., Krieger, M.A., Souto-Padrón, T., De Souza, W. and Goldenberg, S.  
Structure and expression of two *Trypanosoma cruzi* genes encoding antigenic proteins bearing repetitive epitopes, 127

Leech, J.H., see Rosenthal, P.J., 177

Linss, J., see Lafaille, J.J., 127

Lopez, M., see Rege, A.A., 89

Maloy, W.L., see Panton, L.J., 149

Maroney, P.A., Hannon, G.J. and Nilsen, T.W.  
Accurate and efficient RNA polymerase III transcription in

a cell-free extract prepared from *Ascaris suum* embryos, 277

Matile, H., see Caspers, P., 185

McGuire, T.C., see Reduker, D.W., 239

McKerrow, J.H., see Rosenthal, P.J., 177

McPhie, P., see Panton, L.J., 149

McReynolds, L.A., see Grandea III, A.G., 31

Meshnick, S.R., see Kwakye-Berko, F., 51

Mitchell, G.F., see Rogers, M.V., 79

Myler, P.J., see Scholler, J.K., 11

Nilsen, T.W., see Maroney, P.A., 277

Ordóñez, D., see Balaña-Fouce, R., 43

Panton, L.J., McPhie, P., Maloy, W.L., Wellems, T.E., Taylor, D.W. and Howard, R.J.

Purification and partial characterization of an unusual protein of *Plasmodium falciparum*: histidine-rich protein II, 149

Parkhouse, R.M.E., see Harnett, W., 119

Pascal Jr., R.A., see Beach, D.H., 57

Pearson, T.W., see Tolson, D.L., 109

Perryman, L.E., see Reduker, D.W., 239

Philipp, M., see Grandea III, A.G., 31

Piessens, W.F., see Fuhrman, J.A., 249

Pink, J.R., see Caspers, P., 185

Rajan, T.V., see Werner, C., 209

Rasnick, D., see Rosenthal, P.J., 177

Ray, S., see Ghoshal, K., 21

Reduker, D.W., Jasmer, D.P., Goff, W.L., Perryman, L.E., Davis, W.C. and McGuire, T.C.

A recombinant surface protein of *Babesia bovis* elicits bovine antibodies that react with live merozoites, 239

Rege, A.A., Herrera, P.R., Lopez, M. and Dresden, M.H.

Isolation and characterization of a cysteine proteinase from *Fasciola hepatica* adult worms, 89

Remold, H., see Toye, P., 1

Renz, A., see Harnett, W., 119

Rogers, M.V., Henkle, K.J., Fidge, N.H. and Mitchell, G.F.

Identification of multispecific lipoprotein receptor in adult *Schistosoma japonicum* by ligand blotting analyses, 79

Rosenthal, P.J., McKerrow, J.H., Rasnick, D. and Leech, J.H.

*Plasmodium falciparum*: inhibitors of lysosomal cysteine proteinases inhibit a trophozoite proteinase and block parasite development, 177

Schoenmakers, J.G.G., see Wesseling, J.G., 167

Scholler, J.J., Myler, P.J. and Stuart, K.D.

A novel telomeric gene conversion in *Trypanosoma brucei*, 11

Sherman, I.W., see Zidovetzki, R., 199

Silva Filho, F.C., Breier-Saraiva, E.M., Tosta, M.X. and De Souza, W.

*Trichomonas vaginalis* and *Tritrichomonas foetus* secrete neuraminidase into the culture medium, 73

Sinigaglia, F., see Caspers, P., 185

Smits, M.A., see Wesseling, J.G., 167

Snijders, P.J.F., see Wesseling, J.G., 167

Sohal, S., see Allsopp, B., 137

Souto-Padrón, T., see Lafaille, J.J., 127

Stuart, K.D., see Scholler, J.K., 11

Suplick, K., see Vaidya, A.B., 97

Swamy, K.H.S.

Filarial parasites exhibit unusually high levels of choline acetyltransferase activity, 259

Taylor, D.W., see Panton, L.J., 149

Tolson, D.L., Turco, S.J., Beecroft, R.P. and Pearson, T.W.

The immunochemical structure and surface arrangement of *Leishmania donovani* lipophosphoglycan determined using monoclonal antibodies, 109

Tosta, M.X., see Silva Filho, F.C., 73

Toye, P. and Remold, H.

The influence of temperature and serum deprivation on the synthesis of heat-shock proteins and alpha and beta tubulin in promastigotes of *Leishmania major*, 1

Turco, S.J., see Tolson, D.L., 109

Tuyen, L.K., see Grandea III, A.G., 31

Vaidya, A.B., Akella, R. and Suplick, K.

Sequences similar to genes for two mitochondrial proteins and portions of ribosomal RNA in tandemly arrayed 6-kilobase-pair DNA of a malarial parasite, 97

Van Someren, P., see Wesseling, J.G., 167

Wang, C.C., see Hedstrom, L., 219

Wellems, T.E., see Panton, L.J., 149

Werner, C., Higashi, G.I., Yates, J.A. and Rajan, T.V.

Differential recognition of two cloned *Brugia malayi* antigens by antibody class, 209

Wesseling, J.G., Snijders, P.J.F., Van Someren, P., Jansen, J., Smits, M.A. and Schoenmakers, J.G.G.

Stage-specific expression and genomic organization of the actin genes of the malaria parasite *Plasmodium falciparum*, 167

Yates, J.A., see Werner, C., 209

Zidovetzki, R., Sherman, I.W., Atiya, A. and De Boeck, H.

A nuclear magnetic resonance study of the interactions of the antimalarials chloroquine, quinacrine, quinine and mefloquine with dipalmitoylphosphatidylcholine bilayers, 199

## Subject Index

Acetylcholine, 259  
 Actin gene, 167  
 Adenosine, 269  
 Affinity-purified antibody, 31  
 Altered nucleoside permeation, 269  
 Antigen, 127  
 Antigenic variation, 11  
 Antimalarial, 51  
 Antimalarial drug, 149  
 Apoprotein, 79  
*Ascaris suum*, 191, 277  
  
*Babesia bovis*, 239, 269  
 Berenil, 43  
 Bovine erythrocyte, 269  
*Brugia malayi*, 209, 249, 259, 277  
*Brugia pahangi*, 259  
  
 Calcium-binding protein, 249  
 Cell invasion, 229  
 Chagas' disease, 127  
 Chemotherapy, 177  
 Chloroquine, 51  
 Choline acetyltransferase, 259  
 Circumsporozoite protein gene, 185  
 Collagen, 209  
 Cysteine proteinase, 89  
 Cytochrome *b*, 97  
 Cytochrome oxidase, 97  
 Cytotoxicity, 73  
  
 $\alpha$ -Difluoromethylornithine, 43  
 Dihydrosterculinic acid, 47  
*Dirofilaria immitis*, 31  
 DNA, 51  
 rDNA, 67  
 DNA probe, 119  
  
 Ethacrynic acid, 259  
 Evolution, 97  
  
*Fasciola hepatica*, 89, 161  
 Fatty acyl group, 57  
 Filariae, 259  
 Fluoromethylketone, 177  
 Fusion protein, 239  
  
 Gene cloning, 127  
 Gene conversion, 11  
 Glycoprotein, 229  
  
 Haemoglobin, 177  
 Heat-shock protein, 1  
 Histidine-rich protein, 149  
 Hypoxanthine, 219  
  
 IgE, 209  
 IgG, 209  
 In vitro released product, 161  
 In vitro transcription, 277  
 Intercalator, 51  
 Intergenic spacer, 67  
 Internal transcribed spacer, 67  
 Intestine, 191  
 Intron, 167  
  
 D-Lactate, 21  
*Leishmania*, 57  
*Leishmania donovani*, 21, 109  
*Leishmania donovani infantum*, 43  
*Leishmania infantum*, 43  
*Leishmania major*, 1, 109  
 Ligand blotting, 79  
 Lipid membrane, 199  
 Lipophosphoglycan, 109  
 Lipoprotein receptor, 79  
 [ $^{125}$ I]Lipoprotein, 79  
*Litomosoides carinii*, 259  
 Lymphatic filariasis, 209, 259  
 $\lambda$ gt11 cDNA library, 31  
 $\lambda$ gt11 fusion protein expression, 209  
 $\lambda$ gt11 genomic library, 239  
  
 Malaria, 51, 177, 185  
 Methylglyoxal, 21  
 Microfilariae, 249  
 Mitochondrial DNA, 97  
 Monoclonal antibody, 109, 239, 249  
 Muscle, 191  
 Mycophenolic acid resistance, 219  
 Myosin, 31, 209  
  
 Neuraminidase, 73  
 Nitrobenzylthioinosine, 269  
 Non-coding region, 167  
 Nuclear magnetic resonance, 199  
 Nucleoside transport, 269  
  
 Oligonucleotide probe, 137  
*Onchocerca volvulus*, 119  
  
 Paramyosin, 31  
 Parasitic nematode, 191  
 Parasitic protozoon, 73  
 Phosphatidylcholine, 57  
 Phosphatidylethanolamine, 57  
*Plasmodium*, 51, 97  
*Plasmodium falciparum*, 149, 167, 177, 185, 269  
 Polyamine, 43  
 Polymerase chain reaction, 137  
 Promastigote, 57, 109

Protein structure, 149  
Protease inhibitor, 161  
Proteinase, 177  
Protozoa, 21  
Purine transport, 219  
Putrescine transport, 43  
Recombinant protein, 239  
Repeat unit, 67  
Repetitive epitope, 127  
Restriction mapping, 67  
Ribosomal RNA, 97  
RNA polymerase III, 277  
5S rRNA, 277  
*Schistosoma japonicum*, 79  
*Schistosoma mansoni*, 79  
Serotonin receptor, 191  
Speciation, 119  
Stage-specific transcription, 167  
Strain characterization, 137  
Surface epitope, 109  
Surface heterogeneity, 229  
Surface-exposed epitope, 239  
Telomere, 11  
*Theileria parva*, 137  
Thiostearic acid, 57  
Thiol protease, 161  
Transmission-blocking immunity, 249  
Trematode, 161  
Trematode, parasitic, 89  
*Trichinella spiralis*, 67  
*Trichomonas vaginalis*, 73  
*Tritrichomonas foetus*, 73, 219  
*Trypanosoma brucei*, 11  
*Trypanosoma cruzi*, 127, 229  
Tubulin, 1  
Worm motility, 259  
*Wuchereria bancrofti*, 259  
Zinc-chelate chromatography, 149

